Fare Game: A Mean Field Model of Stochastic Intensity Control in Dynamic Ticket Pricing

Burak Aydın * Emre Parmaksız [†] R

Ronnie Sircar[‡]

May 7, 2025

Abstract

We study the dynamic pricing of discrete goods over a finite selling horizon. One way to capture both the elastic and stochastic reaction of purchases to price is through a model where sellers control the intensity of a counting process, representing the number of sales thus far. The intensity describes the probabilistic likelihood of a sale, and is a decreasing function of the price a seller sets. A classical model for ticket pricing, which assumes a single seller and infinite time horizon, is by Gallego and van Ryzin (1994) and it has been widely utilized by airlines, for instance. Extending to more realistic settings where there are multiple sellers, with finite inventories, in competition over a finite time horizon is more complicated both mathematically and computationally. We introduce a dynamic mean field game of this type, and some numerical and existence results. In particular, we analyze the associated coupled system of Hamilton-Jacobi-Bellman and Kolmogorov differential-difference equations, and we prove the existence and uniqueness results under certain conditions. Then, we demonstrate a numerical algorithm to find this solution and provide some insights into the macroeconomic market parameters. Finally, we present a preliminary comparison of our findings with airfare data.

1 Introduction

Deregulation of the airline industry in 1978 ignited the field of revenue management, which was initially known as yield management. Faced with new competitive dynamics, airlines adopted inventory control and dynamic pricing to optimize revenue. These techniques have since diffused across numerous industries, including transportation, entertainment, hospitality, and retail. This paper proposes dynamic pricing strategies for discrete goods such as tickets for an event, in a competitive market over a finite time horizon corresponding to the date of the event.

Sales are modeled as Poisson-type random arrivals of customers to a representative seller. The probability (or intensity) of these arrivals is decreasing in the individual seller's price,

^{*}Department of Operations Research & Financial Engineering, Princeton University, ba7735@princeton.edu

[†]Department of Mathematics, Massachusetts Institute of Technology, *yep@mit.edu*

[‡]Department of Operations Research & Financial Engineering, Princeton University, *sircar@princeton.edu*

and it is increasing with the average market price at any given time. Finite player nonzerosum stochastic differential games of this type are analytically challenging, even with just two firms, as shown in [Ledvina (2011), 21]. They become computationally intractable with more than five firms. To obtain a computationally feasible approximation that yields interpretable results, we propose a mean field game model, which assumes a continuum of firms across the inventory space.

A pioneering work by [Belobaba (1987), 1] underscored how initial airline seat inventory practices relied heavily on human judgment to balance discounted and full-fare tickets for revenue maximization. Recognizing the need for systematic approaches, the Expected Marginal Seat Revenue model was subsequently introduced [Belobaba (1989), 2], which significantly advanced revenue management by incorporating probabilistic decision-making for optimal booking limits. These analytical methodologies demonstrated substantial economic benefits. With advances in computational power, dynamic pricing grew prominent across industries, as extensively reviewed by [McGill and van Ryzin (1999), 24] in the areas of forecasting, overbooking, seat inventory control, and pricing.

Dynamic pricing and inventory management under stochastic demand were rigorously formalized in [Gallego and van Ryzin (1994), 15], where they consider a firm operating in a market with imperfect competition, e.g. a monopoly, such that the firm has the ability to influence its demand by changing the price of a product. The demand in their model is given by a Poisson-type process with intensity $\lambda(p)$, which is non-increasing in price p. At every arrival, one unit is sold at the quoted price, and inventory is decreased by one, until it reaches zero. Their framework illustrated the efficacy of dynamic pricing policies.

A related direction is discrete-choice models introduced by [Talluri and van Ryzin (2004), 25], which explicitly incorporated consumer substitution behavior, allowing more realistic and profitable pricing strategies. Inventory considerations within dynamic pricing contexts were extensively reviewed by [Elmaghraby and Keskinocak (2003), 12], providing critical insights applicable across airlines, hospitality, and retail sectors. This broadened the applicability of dynamic pricing methodologies beyond their traditional scopes. A more recent survey on dynamic pricing (and learning) can be found in [den Boer (2015), 10].

Recent research on oligopolistic airline markets, such as [Betancourt et al. (2022), 3], extended earlier monopolistic frameworks to duopolies and multi-firm setups, highlighting strategic interactions under dynamic pricing with finite inventory and selling horizons. They manage to show improvements in output at the expense of lower welfare levels due to dynamic pricing. Integrating machine learning and game theory into revenue management has also become a prominent area of development. Various methods of game-theory-based reinforcement learning in oligopolistic contexts are analyzed in [Collins and Thomas (2012), 9], and comparisons between reinforcement learning and data-driven programming are discussed in [Lange et al. (2025), 20]. However, as shown in [Bondoux et al. (2020), 4], although these reinforcement learning models may be proven to converge to the optimal solution in the market, in practice they require very large amounts of data. Due to limitations with scarce data, we instead introduce a model-based game theoretic approach.

Another branch of literature that is of interest to our paper is stochastic intensity control, which has also gained considerable attention parallel to revenue management research. Recognizing the need for stochastic intensity control models in perishable goods pricing, earlier models were generalized to allow nonhomogeneous but deterministic Poisson consumer arrival rates in [Zhao and Zheng (2000), 28], significantly improving revenue outcomes compared to static approaches. Competitive revenue management under deterministic demand arrival rates was analyzed by [Gallego and Hu (2014), 14], who studied dynamic pricing equilibrium through shadow prices and extended results to stochastic intensity-controlled games. Similarly, intensity-controlled stochastic exploration in dynamic Cournot games was studied in [Ludkovski and Sircar (2012), 23], revealing strategic adjustments firms make under uncertainty to manage exhaustible resources. These insights are applicable to stochastic demand pricing scenarios.

Despite extensive studies on revenue management, intensity control, and their intersections, mean-field games of intensity control for markets characterized by nonhomogeneous consumer arrival rates remain unexplored. Our paper fills this gap, providing a novel analysis of competition through mean-field game theory as an alternative to analytically and numerically challenging finite-horizon multi-firm competitive models. Mean-field game (MFG) theory is an active research area, thoroughly covered in texts such as [Carmona and Delarue (2018), 6]. It has been applied to various problems related to ours, including Bertrand and Cournot mean-field games [Chan and Sircar (2015), 7], optimal exploitation of exhaustible resources [Graewe et al. (2022), 19], oil markets and fracking [Chan and Sircar (2017), 8], optimal execution in a dynamic pricing context [Donnelly and Li (2022), 11], portfolio liquidation problems [Fu et al. (2021), 13], or cryptocurrency mining [Li et al. (2024), 22]. In the context of the latter model, we refer also to the recent existence results for MFGs of intensity control in [García et al. (2025), 16].

The remainder of the paper is organized as follows. Section 2 introduces the detailed model and its associated coupled Hamilton-Jacobi-Bellman and Kolmogorov differentialdifference equations that characterize the mean field market equilibrium. Our main existence and uniqueness theorems are given in Section 2.3, with their proofs in Appendix C. Section 3 outlines the numerical algorithm, shows our results, discusses the algorithm's numerical stability and effectiveness, and presents a preliminary comparison of our model with airfare data. Finally, Section 4 concludes and discusses directions for future research.

2 Stochastic Intensity Mean Field Game

We consider a continuum of firms distributed across our state space, comprising positive integers. The states represent the number of remaining items, such as tickets, in each firm's inventory. The model encompasses two primary interactions. First, we have the individual optimization of the firms in response to average market prices, strategically adjusting their prices based on their competitors' actions. Second, we have the dynamics of the market itself, which are shaped by the pricing strategies adopted by each firm. The general market dynamics and the actions of individual terms are interdependent through the customer arrival intensity function. A mean field equilibrium is reached when the average of individual firm best responses to a posited mean price, coincides with that mean price (at every time). In other words there is an optimization coupled with a fixed point.

2.1 Model Dynamics

We denote the distribution of firms' inventories by $m(t) = \{m_k(t)\}_{k=0}^K$, where $m_k(t)$ is the proportion of the firms with k items in their inventory at time t, and $K < \infty$ denotes the maximum possible inventory level. We have $\sum_{k=0}^{K} m_k(t) = 1$ for all times $0 \le t \le T$, where $T < \infty$ is the event horizon (for instance time of flight or concert). Firms have finite inventories and are incentivized to sell, as any unsold inventory holds no residual value at the end of the selling period. The initial inventory distribution is given by $m_k(0) = M_k$.

If we denote a firm's inventory by X, its dynamics are

$$dX_t = -\mathbb{1}_{\{X_t > 0\}} dN_t^\lambda,$$

where N_t^{λ} is a counting process with stochastic and time-varying intensity $\lambda_t \geq 0$, meaning

$$\mathbb{P}(N_{t+h}^{\lambda} - N_t^{\lambda} = 1) = \lambda_t h + o(h), \qquad \mathbb{P}(N_{t+h}^{\lambda} - N_t^{\lambda} \ge 2) = o(h).$$

See for instance [Brémaud (1981), 5]. The intensity λ_t is a function that depends on the price p_t quoted by the firm, the average price $\bar{p}(t)$ in the market, and the firm inventory distribution m on the state space:

$$\lambda_t = \lambda(p_t, \bar{p}(t), m(t)).$$

The function λ is assumed decreasing in the firm's own price p_t , so decreasing price increases the probability of a sale. It is increasing in the average price in $\bar{p}(t)$, so with a higher average price, a firm will expect more demand.

A representative seller optimizes expected discounted revenue and the value functions $V(t) = \{V_k(t)\}_{k=0}^K$ when they have k tickets at time $t \leq T$ for each k are given by

$$V_k(t) = \sup_{p_s \ge 0} \mathbb{E} \left\{ \int_t^T e^{-r(s-t)} p_s \mathbb{1}_{\{X_s > 0\}} dN_s^{\lambda} \mid X_t = k \right\},\$$

where the supremum is taken over adapted price trajectories p_s over the trading period $t \leq s \leq T$.

For $k \in \mathcal{K} := \{1, \ldots, K\}$, the associated HJB differential-difference equation is

$$\frac{dV_k}{dt} - rV_k + \sup_{p \ge 0} \left\{ \lambda(p, \bar{p}, m)(p - \Delta V_k) \right\} = 0, \qquad (2.1)$$

where we define the discrete difference $\Delta V_k := V_k - V_{k-1}$. We have the terminal and boundary conditions

$$V_k(T) = 0, \quad k \in \mathcal{K}, \qquad V_0(t) = 0, \quad t \in [0, T],$$

where the first condition means that the remaining goods in the inventory at the end of the selling period will remain unsold and thus generate no value. The second condition means for a firm starting the trading period with zero inventory, the value function is identically zero.

For simplicity, we will consider smooth $\lambda(p, \bar{p}, m)$, and assume that there is a unique supremum in (2.1), which is expressed as

$$p_k^* = \operatorname*{argmax}_{p \ge 0} \Big\{ \lambda(p, \bar{p}, m)(p - \Delta V_k) \Big\}.$$

We define the mean-field equilibrium average price in the market at time t as:

$$\bar{p}(t) = \frac{1}{\eta(t;m)} \sum_{k=1}^{K} m_k(t) p_k^*(t),$$

where

$$\eta(t;m) := \sum_{k=1}^{K} m_k(t) = 1 - m_0(t)$$

is the proportion of active firms. It is important to note that the inactive sold-out firms are omitted from the average calculation, as $\{k = 0\}$ is an absorbing state. We need to keep track of the inactive firms as they do not affect the average price.

The second main equation for our system, known as the Kolmogorov equation, describes the time-evolution of the distribution m(t) when the firms set prices $\{p_k^*(t)\}_{k=1}^K$:

$$\frac{dm_0}{dt} = \lambda(p_1^*, \bar{p}, m) m_1(t),$$

$$\frac{dm_k}{dt} = \lambda(p_{k+1}^*, \bar{p}, m) m_{k+1}(t) - \lambda(p_k^*, \bar{p}, m) m_k(t), \qquad k \in \mathcal{K}_{-1}$$

$$\frac{dm_K}{dt} = -\lambda(p_K^*, \bar{p}, m) m_K(t).$$
(2.2)

subject to an initial condition $m_k(0) = M_k$, and $\mathcal{K}_{-1} := \{1, \ldots, K-1\}$. For example, the second line of (2.2) describes how m_k increases if a firm with inventory k + 1 receives a demand, therefore its inventory level reduces to k, or how it decreases if a k-inventory firm receives a demand and drops to k - 1 inventory.

The HJB and Kolmogorov equations together characterize the mean field equilibrium in the market. Their coupling through the intensity function $\lambda(p, \bar{p}, m)$ allows us to analyze the interplay between firm strategies, market dynamics, and the resulting equilibrium as the system evolves over time.

2.2 Linear Intensity Model

In the remainder of the paper, we work with the linear intensity function

$$\lambda(p,\overline{p},m) = (a(m) - p + c(m)\overline{p})^+, \qquad (2.3)$$

where

$$a(m) = \frac{1}{1 + \epsilon \sum_{k=1}^{K} m_k} = \frac{1}{1 + \epsilon \eta(m)}, \qquad c(m) = \frac{\epsilon \eta(m)}{1 + \epsilon \eta(m)},$$

and we have suppressed the *t*-dependence of m and η in the notation. We observe that a(m) = 1 - c(m). The parameter $\epsilon \ge 0$ represents the level of competition in the market. When $\epsilon = 0$, $\lambda(p, \overline{p}, m)$ becomes independent of m and \overline{p} and the firms behave like monopolies.

The function in (2.3) satisfies $\partial_p \lambda \leq 0$ and $\partial_{\overline{p}} \lambda \geq 0$. Moreover, when all firms are inactive $(m_0 = 1, m_k = 0 \ \forall k \in \mathcal{K})$, we have $\eta(m) = 0$, and we observe that $\lambda(p, \overline{p}, m) = (1-p)^+$ depends only on p. This shows how a firm becomes closer to a monopoly as the other firms

exit the market by exhausting their inventories. Our motivation to consider the linear intensity function comes from its derivation in [Chan and Sircar (2015), 7]. Referring to [Vives (2001), 26, Chapter 6], they consider finitely many agents that have a linear price-demand relation, and by solving the system of equations relating to the price-demand equilibrium, they obtain a relation of the form (2.3) in the continuum limit $N \to \infty$. A brief discussion is given in Appendix A.

Using the expression for λ in (2.3), and suppressing a and c's dependence on m, we obtain

$$p_k^* = \frac{1}{2} \left(a + c\overline{p} + \Delta V_k \right)^+. \tag{2.4}$$

As demonstrated in the proof our main theorems, given in Appendix C, the quantity between the parentheses in (2.4) is non-negative in equilibrium for small enough ϵ , so we can remove the positive part. Then we obtain a simplified expression for the average price \bar{p} :

$$\overline{p} = \frac{1}{2-c} \left(a + \frac{1}{\eta} \sum_{k=1}^{K} m_k \Delta V_k \right).$$

Plugging these expressions for p_k^* and \bar{p} into (2.1) and (2.2), and defining the function

$$\phi_{\epsilon}(V,m) := \frac{1}{2 + \epsilon \eta(m)} \Big(\sum_{k=1}^{K} m_k \Delta V_k - \eta(m) \Big),$$

we arrive at the coupled system of differential equations that together govern the dynamics of the model:

$$\frac{dV_k}{dt} - rV_k + \frac{1}{4} \left(1 - \Delta V_k + \epsilon \phi_\epsilon(V, m) \right)^2 = 0, \quad k \in \mathcal{K}$$

$$\frac{dm_k}{dt} - \left[\frac{1}{2} \left(1 - \Delta V_{k+1} + \epsilon \phi_\epsilon(V, m) \right) m_{k+1} - \frac{1}{2} \left(1 - \Delta V_k + \epsilon \phi_\epsilon(V, m) \right) m_k \right] = 0, \quad k \in \mathcal{K}_{-1}$$

$$\frac{dm_0}{dt} - \frac{1}{2} \left(1 - \Delta V_1 + \epsilon \phi_\epsilon(V, m) \right) m_1 = 0,$$

$$\frac{dm_K}{dt} + \frac{1}{2} \left(1 - \Delta V_K + \epsilon \phi_\epsilon(V, m) \right) m_K = 0,$$

$$V_k(T) = 0, \quad k \in \mathcal{K}, \quad m_k(0) = M_k, \quad k \in \mathcal{K}.$$
(2.5)

A direct observation shows that when the interaction parameter ϵ is identically zero, the two main equations relating V and m are decoupled. We argue that in the case of non-zero but sufficiently small competition ($\epsilon > 0$), we can prove existence and uniqueness by linearizing the system around the monopoly case and using the Schauder Fixed Point Theorem [Gilbarg and Trudinger (2001), 17, Corollary 11.2, p. 280]. Our proofs are related to those for the continuous-space Cournot exhaustible resources oil extraction MFG model in [Graber and Sircar (2023), 18].

2.3 Main Results

The following proposition will be useful in the proofs of the main theorems.

Proposition 2.1. For $\epsilon = 0$, a solution $(V^{(0)}, m^{(0)}) = (\{V_k^{(0)}\}_{k=1}^K, \{m_k^{(0)}\}_{k=1}^K)$ to the system of differential equations (2.5) exists and is unique. Further, for all $k \in \mathcal{K}$, and $t \in [0, T]$, we have

$$0 \leq \Delta V_k^{(0)}(t) \leq 1 - 2\rho < 1$$

$$\frac{d\Delta V_k^{(0)}}{dt}(t) \leq 0$$

$$0 \leq \eta^{(0)}(t; m^{(0)}) \leq 1$$

$$0 \leq m_k^{(0)}(t) \leq 1,$$

(2.6)

where $\rho = \sqrt{r^2 + r} - r \in [0, \frac{1}{2})$. Consequently, for $\lambda_k^{(0)}(t) = \frac{1}{2}(1 - \Delta V_k^{(0)}(t))$, we have

$$\rho \le \lambda_k^{(0)}(t) \le \frac{1}{2}.$$

The proof is given in Appendix B. The bounds (2.6) are in line with our expectations for a solution that is economically sensible. The parameter $\lambda_k^{(0)}$ represents customer arrival rates, hence it should be non-negative as depicted. Similarly, the variables m_k and η signify proportions, and thus it is crucial for these values to lie within the range [0, 1] to align with economic rationale. A noteworthy detail is that the $\lambda_k^{(0)}$ has a lower bound of $\rho > 0$, uniformly across k and in t.

Our two main theorems are the following:

Theorem 2.2. In the setting described above, for any initial inventory distribution $\{M_k\}_{0 \le k \le K}$, interest rate r > 0 and time horizon T > 0, there exists a solution $\{V_k(t), m_k(t)\}_{0 \le k \le K}$ to the HJB-Kolmogorov system of equations if the interaction parameter $\epsilon > 0$ is less than some constant C_1 , which depends on the model parameters T, K, r.

Theorem 2.3. In the setting described above, for any initial inventory distribution $\{M_k\}_{0 \le k \le K}$, interest rate r > 0 and time horizon T > 0, there exists an upper bound $C_2 < C_1$ such that if $\epsilon < C_2$, then the solution exists and is unique.

The proofs of these theorems are in Appendix C. For the proof of existence, we consider first-order deviations of the solution (V, m) to (2.5), from the monopolist solution $(V^{(0)}, m^{(0)})$, prescribe a compact set as their domain M, and derive the differential equations that control them. Deriving the integral operators whose fixed points give the solutions to the differential equations, we apply the Schauder Fixed Point Theorem after checking for the operator properties such as mapping M into itself, continuity, and precompactness of the image of M. For uniqueness, we utilize an energy identity. Assuming two solutions to (2.5), we prove under restrictions on ϵ that the solutions are indeed the same.

These theorems highlight the existence and structure of equilibrium solutions, laying a solid foundation for subsequent numerical investigations in Section 3.

3 Numerical Results

In this section, our primary goal is to establish a numerical method for computing the meanfield equilibrium. Subsequently, we will demonstrate the convergence of our theoretical findings to the numerical results in the limit $\epsilon \to 0$. Finally, we will conclude this section by conducting an in-depth analysis of the numerical stability of our solution strategy and explore various ways to establish theoretical convergence proofs.

3.1 Algorithm

As discussed in Section 2, the mean-field equilibrium characterizes a state in which the aggregate behavior of the continuous set of players reaches a fixed point of an optimization problem. In this equilibrium, each player autonomously adopts a Markovian strategy to optimize their individual value function while taking into account the collective behavior of the entire population. As a result, these individually optimized strategies jointly contribute to the overall movement of the population as initially assumed. Our algorithm for finding this fixed point is as follows:

Step 0: Initial guess

Begin with an initial guess $(\bar{p}^{(0)}(t), m^{(0)}(t))$, and for n = 0, 1, 2, ... do the following steps:

Step 1: Solving the HJB equation

We discretize the HJB equation in time. Using the initial guess $(\bar{p}^{(n)}(t), m^{(n)}(t))$, we employ the Runge-Kutta method for ODEs of order 5(4) to solve.

$$\frac{dV_k}{dt} - rV_k + \sup_p \left\{ \lambda(p, \bar{p}^{(n)}(t), m^{(n)}(t))(p - \Delta V_k) \right\} = 0, \quad k \in \mathcal{K}$$

with the terminal condition $V_k(T) = 0$. Using the solution, find the prices $p_k^{*(n)}(t)$ and intensities $\lambda_k^{*(n)}(t) := \lambda(p_k^{*(n)}(t), \bar{p}^{(n)}(t), m^{(n)}(t))$.

Step 2: Solving the Kolmogorov equation

Using intensities in step 1, find $m^{(n+1)}(t)$ by solving the Kolmogorov equation:

$$\frac{dm_0^{(n+1)}}{dt} = \lambda_1^{*(n)}(t)m_1^{(n+1)}
\frac{dm_k^{(n+1)}}{dt} = -\lambda_k^{*(n)}(t)m_k^{(n+1)} + \lambda_{k+1}^{*(n)}(t)m_{k+1}^{(n+1)}, \qquad k \in \mathcal{K}_{-1}
\frac{dm_K^{(n+1)}}{dt} = -\lambda_K^{*(n)}(t)m_K^{(n+1)}$$

similarly. Using the solution, find $\bar{p}^{(n+1)}(t)$:

$$\bar{p}^{(n+1)}(t) = \frac{\sum_{k=1}^{K} m_k^{(n+1)}(t) p_k^{*(n)}(t)}{\sum_{k=1}^{K} m_k^{(n+1)}}$$

If $(\bar{p}^{(n+1)}(t), m^{(n+1)}(t))$ and $(\bar{p}^{(n)}(t), m^{(n)}(t))$ are sufficiently close to each other, we terminate. Otherwise, we go back to Step 1 with the newly computed $(\bar{p}^{(n+1)}(t), m^{(n+1)}(t))$.

Remarks on the algorithm

• As a proxy for distance between $(\bar{p}^{(1)}(t), m^{(1)}(t))$ and $(\bar{p}^{(2)}(t), m^{(2)}(t))$, we take

$$L_{\bar{p}}(\bar{p}^{(1)}, \bar{p}^{(2)}) := \int_{0}^{T} \left| \bar{p}^{(1)}(t) - \bar{p}^{(2)}(t) \right|^{2} dt,$$
$$L_{m}(m^{(1)}, m^{(2)}) := \sum_{k=0}^{K} \int_{0}^{T} \left| m_{k}^{(1)}(t) - m_{k}^{(2)}(t) \right|^{2} dt,$$
$$L((\bar{p}^{(1)}, m^{(1)}), (\bar{p}^{(2)}, m^{(2)})) := L_{\bar{p}}(\bar{p}^{(1)}, \bar{p}^{(2)}) + L_{m}(m^{(1)}, m^{(2)})$$

• We calculate the time integrals for the error term using the same partition as we had while solving HJB and Kolmogorov equations.

3.2 Numerical Stability and Convergence of the Algorithm

To test the numerical stability of the algorithm and the uniqueness of the equilibrium point, we simulate a sample system with N-many random initial guesses and check the corresponding value functions $\{V_k^{(1)}(t), \ldots, V_k^{(N)}(t)\}$, price profiles $\{\bar{p}_k^{(1)}(t), \ldots, \bar{p}_k^{(N)}(t)\}$ and inventory distributions $\{m_k^{(1)}(t), \ldots, m_k^{(N)}(t)\}$.

For each quantity, for each time point t, and for each level k > 0, we calculate the "coefficient of variation" (standard deviation / mean) from N samples and finally, take the supremum of this quantity over the entire period, among all the levels.

Mathematically, for a generic quantity $\{q_k(t)\}_{k=1,t=0}^{K,T}$, we use the following formula:

$$CV_q = \sup_{k \in \mathcal{K}, t \in [0,T]} \frac{\sqrt{\frac{1}{N} \left[\sum_{n=1}^{N} \left(q_k^{(n)}(t) - \frac{1}{N} \sum_{m=1}^{N} q_k^{(m)}(t) \right)^2 \right]}}{\frac{1}{N} \sum_{n=1}^{N} q_k^{(n)}(t)}.$$

For the values N = 10, T = 200, r = 0.04, $\epsilon = 0.4$, we find

$$CV_V \approx 4 \times 10^{-11}, \ CV_{\bar{p}} \approx 5 \times 10^{-12}, \ CV_m \approx 2 \times 10^{-8}.$$

Based on our observations, the convergence is typically achieved in 6 steps, and the proportional decrease in error remains consistent after the first step. Notably, the proportionality constant changes with ϵ , with smaller values of ϵ leading to faster convergence. This aligns with expectations, as for $\epsilon = 0$, convergence occurs in a single iteration due to the independence of the HJB solution from m(t), $\bar{p}(t)$.

3.3 Algorithm Results

We set K = 100, T = 200 days, r = 0.04 per day. For the initial distribution, we give a "bimodal-like" distribution, comprising two groups of firms uniformly distributed over $k \in \{20, \ldots, 24\} \cup \{50, \ldots, 54\}$, that is, a low-inventory and a high-inventory group. For our initial condition on m, we get the following graphs which shows the concavity of V in k at the starting time t = 0, by demonstrating diminishing marginal returns principle in action: We see that incremental gains from additional inventory reduce as inventory level increases, due to saturation and increased urgency to sell before expiration.



Figure 1: Concavity of V in k

In Figure 2, we plot the cumulative distribution function (CDF) of m at three time instances over inventory level k, for $\epsilon \in \{0, 0.4\}$. The plots show that with increased competition, the market is slower, that is, the firms sell at a slower rate (evidenced by lagging densities when $\epsilon = 0.4$). As firms get more competitive, we get a more stable inventory flow. This suggests that intensified competition leads firms to behave cautiously, pricing tickets more conservatively to avoid rapid inventory depletion and subsequent competitive disadvantage. Hence, increased competition fosters more stable inventory dynamics, ensuring firms maintain availability throughout the selling period.

Figure 3 reinforces this interpretation by displaying optimal price strategies under varying competition levels. Increased competition results in uniformly lower optimal prices across inventory levels, confirming competitive pressures incentivize price reductions to attract customers. Furthermore, higher inventory levels consistently translate to lower prices, indicating urgency in inventory liquidation as the time horizon shortens. This strategic pricing behavior highlights the necessity of balancing revenue generation with the risk of unsold inventory.



Figure 2: Evolution of CDF of m(t) at given times for varying ϵ .



Figure 3: Evolution of optimal price $p^*(t)$ at given times for varying ϵ .



Figure 4: Comparisons for varying ϵ values.

Figure 4 displays the time-evolutions of two important market quantities, proportion of active firms $\eta(t)$ and average price in the market $\bar{p}(t)$ for varying ϵ . The slower rate of selling with higher competition levels is also inferred from the proportion of active firms Figure 4a, as an increased proportion of firms are actively selling inventory over extended periods. In Figure 4b, as expected with more competition, average market price decreases, benefiting consumers. These two effects together underscore competition's role in prolonging market participation, allowing consumers greater purchasing flexibility.

In particular, the inflection points of the $\bar{p}(t)$ curve in Figure 4b correspond to key moments in the evolution of firm density, enhancing the interpretability of the results. The first inflection occurs when the low-inventory group begins to sell out; the second appears as this group approaches complete exit from the market; and the third inflection marks the start of market exit by the high-inventory group.



Figure 5: Density m(t) at inflection points

The model is showing promising results in applicability to real-world data. To see a demonstration on airplane ticket pricing, we used average flight price data (collected on

Tuesdays) for Tuesday morning flights from Chicago O'Hare International Airport (ORD) to New York LaGuardia Airport (LGA), covering five major airlines [Wong (2021), 27]. Our selection criteria focused on flights predominantly taken by business travelers, with the aim of minimizing the price discrimination effects, which are not in the scope of our model.

Figure 6 presents a comparison between the normalized average flight prices on a Tuesday (plots are for September 20, 2022), and the prices predicted by our model. The main difference is the terminal price drop observed in the numerical results but not in the real-world data. This difference can be interpreted to be caused by the fact that pricing is a repeated game with consumers capable of learning in the real world. As a result, firms have little incentive to sharply lower prices in the final days before a flight. On the other hand, our model is a non repeated game in continuous time with no carryover of information, and firms are shown to prefer aggressive end-of-period price reductions to generate some value compared to the zero value of unsold inventories.



Figure 6: Our numerical solution and $ORD \rightarrow LGA$ Tuesday morning flight pricing data

4 Conclusion

In this paper, we introduced a novel existence and uniqueness result for a finite-state mean field game of stochastic intensity control, using differential equations methods. Our approach aims to provide a new perspective on analyzing such games in revenue management, and also to contribute additional tools for studying state mean field interactions in discrete-state settings.

We note that our ticket-selling model can be generalized to encompass a variety of situations seen in the real markets. One of these situations is the option of reselling by the customers. In this case, the price for reselling should also be another control parameter. This can be readily applied in our framework. Another example is the firm overselling. In this instance, firms have the option to oversell their tickets at the cost of paying high penalties at the end of the trading period if they oversell by that time. This example is seen in the airline companies for instance. Our model can be used to understand whether this scheme leads to economically good results for the customers similar to the firm competition. In the computations for the overselling setup, the value functions of the following form can be considered:

$$V_k(t) = \sup_{p_s \ge 0} \mathbb{E} \Big\{ \int_t^T e^{-r(s-t)} p_s \mathbb{1}_{\{X_s > 0\}} dN_s^{\lambda} + e^{-r(T-t)} \Psi(X_T) \Big| X_t = k \Big\},$$

where $\Psi(X_T)$ is a penalty function for overselling, where $\Psi(x) = 0$ for $x \ge 0$, further tailored to suit the desired market model.

We can extend number of tickets being sold at every arrival by considering N as a marked point process. In realistic situations, groups of people traveling together, such as families, would buy their tickets together at the same time, if there is sufficient inventory. Therefore, in our model, each arrival can be associated with a random number of sales, such that the HJB equation becomes

$$\frac{dV_k}{dt} - rV_k + \sup_{p\geq 0} \left\{ \sum_{j=1}^{J\wedge k} \lambda_k \pi_j \left[jp - (V_k - V_{k-j}) \right] \right\} = 0,$$

and Kolmogorov equations can be written as

$$\frac{dm_k}{dt} - \left(\sum_{j=1}^{J \wedge K-k} \pi_j \lambda_{k+j} m_{k+j} - \lambda_k m_k\right) = 0,$$

subject to same initial and terminal conditions as in (2.5).

More general intensity models: as we showed in Appendix C, the existence and uniqueness of a fixed point can be proven for a sufficiently small interaction parameter. Our proof relied upon topological fixed point theorems and an energy identity. These methods are rather canonical, and therefore, the results can be generalized to a larger class of intensity functions. For the sake of brevity, in this paper, we only considered a piecewise-linear case.

We aim to extend our model to incorporate cancellations and refund mechanisms, as experienced in real world due to catastrophes or disruptions such as pandemics, weather anomalies, natural disasters, especially relevant for airline or hospitality industries. Further analysis can also be conducted in the presence of abundant data, both on its own using techniques such as reinforcement learning to approach this problem, and in addition to our method by calibrating the model parameters.

A Brief Review and Limit of the Finite-Player Case

We show that the consumer preferences in our model can be derived from Quasilinear Quadratic Utility Model (QQUM). Letting $q = (q_i)_{i=1}^N$, a suitable utility function is

$$U(q) = \sum_{i=1}^{N} q_i - \frac{1}{2} \left(\sum_{i=1}^{N} q_i^2 + \frac{\epsilon}{N-1} \sum_{j \neq i} q_i q_j \right).$$

The first-order condition of the utility maximization problem $\max_{q \in \mathbb{R}^N_+} U(q) - pq$ for a representative consumer yields the inverse demand functions:

$$q_i = D_i(p) = \begin{cases} a_n - b_n p_i + c_n \bar{p}_i^n, & i = 1, 2, \cdots, n \\ 0, & i = n + 1, \cdots, N \end{cases}$$

where n-many firms receive positive demands, and thus, are active and produce positive quantities, and the rest of the firms are inactive [Vives (2001), 26, see Chapter 6 for more details]. These constants can be calculated as

$$a_n = \frac{1}{1 + \epsilon \frac{n-1}{N-1}}, \quad b_n = \frac{1 + \epsilon \frac{n-2}{N-1}}{(1 + \epsilon \frac{n-1}{N-1})(1 - \frac{\epsilon}{N-1})}, \quad c_n = \frac{\epsilon \frac{n-1}{N-1}}{(1 + \epsilon \frac{n-1}{N-1})(1 - \frac{\epsilon}{N-1})}.$$
 (A.1)

In the context of our paper, for very large (n, N), $\frac{n-1}{N-1} \to \eta(m)$ (the proportion of active firms), and $\frac{1}{N-1} \to 0$. Consequently, the equations in (A.1) become

$$a(m) = \frac{1}{1 + \epsilon \eta(m)}, \qquad b(m) = 1, \qquad c(m) = \frac{\epsilon \eta(m)}{1 + \epsilon \eta(m)},$$

concluding the justification about our demand arrival process rate of

$$\lambda(p,\overline{p},m) = (a(m) + c(m)\overline{p} - p)^+.$$

B Proof of Proposition 2.1

B.1 The inductive setup for Picard-Lindelöf theorem

As the HJB equations become independent from the Kolmogorov equations when $\epsilon = 0$, we shall start by first solving for $V_k^{(0)}$'s, proceeding by induction on k. Remember that the HJB equations are given by

$$\frac{dV_k^{(0)}}{dt} - rV_k^{(0)} + \frac{1}{4}(1 - \Delta V_k^{(0)})^2 = 0.$$
(B.1)

Suppose, for an induction argument, that the bounds in Proposition 2.1 hold for some k. From (B.1), we get

$$\frac{d\Delta V_{k+1}^{(0)}}{dt} + \frac{1}{4} (A_k - \Delta V_{k+1}^{(0)}) (B_k - \Delta V_{k+1}^{(0)}) = 0,$$
(B.2)

where we defined

$$A_k(t) = 1 + 2r - \sqrt{(1+2r)^2 - 1 + (1 - \Delta V_k^{(0)}(t))^2}$$

$$B_k(t) = 1 + 2r + \sqrt{(1+2r)^2 - 1 + (1 - \Delta V_k^{(0)}(t))^2}.$$

Now we argue that a unique solution to (B.2) exists in [0,T] and further it satisfies $0 \leq \Delta V_{k+1}^{(0)}(t) < A_k(t)$ for all $t \in [0,T]$. Introducing a new variable

$$z(t) = \frac{1}{A_k(t) - \Delta V_{k+1}^{(0)}(t)},$$

and $0 < z(T) \le z(t) < \infty$ for all $t \in [0, T]$. Notice that for z, (B.2) can be written as

$$\frac{dz}{dt} + \frac{1}{4}(B_k - A_k)z + A'_k z^2 + \frac{1}{4} = 0.$$
(B.3)

B.2 Inductive step, proving existence and uniqueness of $\Delta V_k^{(0)}$

By Picard-Lindelöf theorem, it suffices to show that the solution does not blow up for any $t_0 \in [0, T]$. Therefore, we will now bound the solution from above and below. We can rewrite (B.3) as

$$z' + 2\alpha z - \beta z^2 + \gamma = 0,$$

where $\alpha(t), \beta(t), \gamma(t)$ are non-negative functions following from $B_k(t) - A_k(t) \ge 0$ and $A'_k(t) \le 0$. Change of variables $\tilde{z}(t) = z(t)e^{\int_0^t \alpha(s)ds}$ yields

$$\tilde{z}' - \tilde{\beta}\tilde{z}^2 + \tilde{\gamma} = 0$$

where $\tilde{\beta}(t) = \beta(t)e^{-\int_0^t \alpha(s)ds} \ge 0$ and $\tilde{\gamma}(t) = \gamma(t)e^{\int_0^t \alpha(s)ds} \ge 0$. Then the upper bound for $\tilde{z}(t)$ can be shown by

$$\tilde{z}(t) \le \tilde{z}(T) + \int_0^T \tilde{\gamma}(s) ds$$

Conversely for the lower bound, inspecting $(1/\tilde{z})' + \tilde{\beta} - \tilde{\gamma}/\tilde{z}^2 = 0$ allows us to show that

$$\tilde{z}(t) \ge \left(\frac{1}{\tilde{z}(T)} + \int_0^T \tilde{\beta}(s)ds\right)^{-1}$$

This concludes the proof that a solution $\Delta V_{k+1}^{(0)}$ in [0, T] exists and is unique. Further for all $t \in [0, T]$

$$0 \le \Delta V_{k+1}^{(0)}(t) \le A_k(t) \le 1 + 2r - 2\sqrt{r^2 + r} = 1 - 2\delta$$

and $\frac{d\Delta V_{k+1}^{(0)}}{dt}(t) \leq 0$, which is observed readily from (B.3) and the fact that $B_k \geq A_k$. This concludes the inductive step. Now let us prove it for the base case.

B.3 Base case, proving existence and uniqueness of $V_k^{(0)}$

In this case, we have

$$\frac{d\Delta V_1^{(0)}}{dt} + \frac{1}{4}(A_1 - \Delta V_1^{(0)})(B_1 - \Delta V_1^{(0)}) = 0,$$

where

$$A_1 = 1 + 2r - 2\sqrt{r^2 + r}$$
$$B_1 = 1 + 2r + 2\sqrt{r^2 + r}.$$

Since A_1 and B_1 have no dependence on t, the solution has an explicit form.

$$\Delta V_1^{(0)}(t) = A_1 \frac{1}{1 + \frac{B_1 - A_1}{A_1} (e^{\sqrt{r^2 + r}(T - t)} - 1)}.$$

It is seen that $\Delta V_1^{(0)}$ satisfies the appropriate conditions. This concludes the proof that there exists a unique solution to $V_k^{(0)}$ in [0, T].

Proving existence and uniqueness of $m_k^{(0)}$ **B.4**

With the results for $V_k^{(0)}$ in mind, recall that the Kolmogorov equations are

$$\frac{dm_k^{(0)}}{dt} - (\lambda_{k+1}^{(0)}m_{k+1}^{(0)} - \lambda_k^{(0)}m_k^{(0)}) = 0, \quad k \in \mathcal{K}_{-1}$$
$$\frac{dm_K^{(0)}}{dt} + \lambda_K^{(0)}m_K^{(0)} = 0,$$

with the initial condition $m_k^{(0)}(0) = M_k$, satisfying $0 \le M_k \le 1, \forall k \in \mathcal{K}$ and $\sum_{k=0}^K M_k = 1$. Consequently, we have the following solutions

$$m_{K}^{(0)}(t) = M_{K} \exp\left(-\int_{0}^{t} \lambda_{K}^{(0)}(s) ds\right)$$
$$m_{k}^{(0)}(t) = \exp\left(-\int_{0}^{t} \lambda_{k}^{(0)}(s) ds\right) \left[M_{k} + \int_{0}^{t} \lambda_{k+1}^{(0)}(s) m_{k+1}^{(0)}(s) \exp\left(\int_{0}^{s} \lambda_{k}^{(0)}(u) du\right) ds\right], \quad k \in \mathcal{K}_{-1}.$$
(B.4)

Since $M_K \ge 0$ and all the other terms are nonnegative in (B.4), by induction, we can see that $\forall k \in \mathcal{K}, \ \forall t \in [0, T], \ m_k^{(0)}(t) \ge 0.$ Further, for $\eta^{(0)}(t; m^{(0)}) = \sum_{k=1}^K m_k^{(0)}(t)$, we have

$$\frac{d\eta^{(0)}}{dt} = -\lambda_1^{(0)} m_1^{(0)},$$

with $\eta^{(0)}(0; m^{(0)}) = 1$ (assuming without loss of generality that no firms start with zero inventory). Therefore $\eta^{(0)}(t; m^{(0)}) \leq 1$ for $t \in [0, T]$, and this implies that for all $t \in [0, T]$

$$0 \le m_k^{(0)}(t) \le 1, \quad k \in \mathcal{K}.$$

This concludes the proof.

C Proofs of Existence and Uniqueness

C.1 Proof of Existence

Let us denote the monopolist solution as $(V^{(0)}, m^{(0)})$, and define the first-order deviations $V^{(1)} = \{V_k^{(1)}\}_{k=1}^K$ and $m^{(1)} = \{m_k^{(1)}\}_{k=1}^K$ by

$$V_k = V_k^{(0)} + \epsilon V_k^{(1)}$$
$$m_k = m_k^{(0)} + \epsilon m_k^{(1)},$$

where $V_k^{(1)}$ and $m_k^{(1)}$ will depend on ϵ . We will consider the vector space of continuous functions

$$\mathcal{B} := \{ (V_1^{(1)}, \dots, V_K^{(1)}, m_1^{(1)}, \dots, m_K^{(1)}), \quad V_k^{(1)}, m_k^{(1)} \in C([0, T]), \quad k \in \mathcal{K} \},\$$

and the uniform norm

$$\|(V^{(1)}, m^{(1)})\|_{\mathcal{B}} = \sum_{k=1}^{K} \|V_k^{(1)}\|_{\infty} + \sum_{k=1}^{K} \|m_k^{(1)}\|_{\infty}$$

With this norm, \mathcal{B} is a Banach space. Vector addition and scalar multiplication are defined elementwise and pointwise as usual.

C.1.1 Outline of the proof process

By formulating two operators R and S, we will transform our problem into finding a fixed point $(V^{(1)}, m^{(1)})$, such that

$$V_k^{(1)} = R_k(V^{(1)}, m^{(1)}), \quad k \in \mathcal{K}$$
$$m_k^{(1)} = S_k(V^{(1)}, m^{(1)}), \quad k \in \mathcal{K}.$$

To use the Schauder Fixed Point Theorem, we will first define the mapping Γ :

$$\Gamma((V^{(1)}, m^{(1)})) = (R_1(V^{(1)}, m^{(1)}), \dots, R_K(V^{(1)}, m^{(1)}), S_1(V^{(1)}, m^{(1)}), \dots, S_K(V^{(1)}, m^{(1)})).$$

Then, we will find the bounds $\{C_k^V, C_k^m\}_{k=1}^K$ such that the following closed and convex ellipsoid

$$M := \{ (V^{(1)}, m^{(1)}) \in \mathcal{B} : \|V_k^{(1)}\|_{\infty} \le C_k^V, \ \|m_k^{(1)}\|_{\infty} \le C_k^m, \ k \in \mathcal{K} \}$$

is mapped into itself under Γ . Thereafter by proving that Γ is continuous and $\Gamma(M)$ is precompact (i.e. its closure is compact), we will prove the existence of a fixed point.

In the proof, we will derive the operators and define the bounds $\{C_k^V\}_{k=1}^K$ and $\{C_k^m\}_{k=1}^K$ as functions solely dependent on the external parameters: T, K, r. Subsequently, we will prove that for a sufficiently small $\epsilon > 0$, the mapping Γ maps M into itself. Next, we will establish the continuity of Γ with respect to the norm $\|\cdot\|_{\mathcal{B}}$. And finally, we will show that the family $\Gamma(M)$ of functions is equicontinuous and pointwise-bounded, which will allow us to conclude that $\Gamma(M)$ is precompact by the Arzela-Ascoli Theorem.

C.1.2 Deriving the integral operators

Our strategy is to linearize (2.5) and solve it using Schauder's Fixed Point Theorem. To begin, let us establish the following notations:

$$\eta^{(0)} = \sum_{k=1}^{K} m_k^{(0)}, \qquad \eta^{(1)} = \sum_{k=1}^{K} m_k^{(1)},$$

$$\phi^{(0)} = \phi_0(V^{(0)}, m^{(0)}) = \frac{1}{2} \Big(\sum_{k=1}^{K} m_k^{(0)} \Delta V_k^{(0)} - \eta^{(0)} \Big).$$
(C.1)

Aiming to satisfy $\phi_{\epsilon}(V^{(1)}, m^{(1)}) = \phi^{(0)} + \epsilon \phi^{(1)}$, we define the first-order perturbation term $\phi^{(1)}$ as

$$\phi^{(1)} = \frac{1}{2 + \epsilon(\eta^{(0)} + \epsilon\eta^{(1)})} \Big(-(\eta^{(0)} + \epsilon\eta^{(1)})\phi^{(0)} + \sum_{k=1}^{K} m_k^{(1)} \Delta V_k^{(0)} + \sum_{k=1}^{K} m_k^{(0)} \Delta V_k^{(1)} - \eta^{(1)} \Big)$$

Then the first equation in (2.5) becomes

$$\frac{dV_k^{(1)}}{dt} - (\lambda_k^{(0)} + r)V_k^{(1)} + \lambda_k^{(0)}(\phi^{(0)} + V_{k-1}^{(1)}) + \epsilon\Lambda_k^V = 0,$$

where $\lambda_k^{(0)}$ and Λ_k^V are defined in the following way:

$$\lambda_k^{(0)} = \frac{1}{2} (1 - \Delta V_k^{(0)})$$

$$\Lambda_k^V = \lambda_k^{(0)} \phi^{(1)} + \frac{1}{4} \left(\phi^{(0)} - \Delta V_k^{(1)} + \epsilon \phi^{(1)} \right)^2.$$

Let us introduce the integral operator

$$R_k(V^{(1)}, m^{(1)})(t) = \int_t^T e^{-\int_t^s (\lambda_k^{(0)}(s') + r)ds'} \left(\lambda_k^{(0)}(\phi^{(0)} + V_{k-1}^{(1)}) + \epsilon \Lambda_k^V\right) ds.$$
(C.2)

With this definition, our problem is transformed into finding a fixed point $V^{(1)}$ under R, that is, for $k \in \mathcal{K}, V_k^{(1)} = R_k(V^{(1)}, m^{(1)})$.

Similarly, let us derive an integral operator for $m^{(1)}$. We start by defining

$$\lambda_k^{(1)} = \frac{1}{2} \Big(\phi^{(0)} - \Delta V_k^{(1)} + \epsilon \phi^{(1)} \Big),$$

the first-order deviation from λ_k such that $\lambda_k \equiv \frac{1}{2}(1 - \Delta V_k + \epsilon \phi_\epsilon) = \lambda_k^{(0)} + \epsilon \lambda_k^{(1)}$. We observe that positivity of λ_k is preserved for small enough $\epsilon > 0$.

We further define

$$\Lambda_k^m = m_{k+1}^{(1)} \lambda_{k+1}^{(1)} - m_k^{(1)} \lambda_k^{(1)}, \text{ for } k \in \mathcal{K}_{-1} \text{ and } \Lambda_K^m = -m_K^{(1)} \lambda_K^{(1)}$$

to simplify the Kolmogorov equations in (2.5) and arrive at the integral operator:

$$S_{k}(V^{(1)}, m^{(1)})(t) = \int_{0}^{t} e^{-\int_{s}^{t} \lambda_{k}^{(0)}(s')ds'} \left(m_{k+1}^{(1)}\lambda_{k+1}^{(0)} + m_{k+1}^{(0)}\lambda_{k+1}^{(1)} - m_{k}^{(0)}\lambda_{k}^{(1)} + \epsilon\Lambda_{k}^{m} \right) ds \quad k \in \mathcal{K}_{-1}$$

$$S_{K}(V^{(1)}, m^{(1)})(t) = \int_{0}^{t} e^{-\int_{s}^{t} \lambda_{k}^{(0)}(s')ds'} \left(-m_{K}^{(0)}\lambda_{K}^{(1)} + \epsilon\Lambda_{K}^{m} \right) ds.$$
(C.3)

C.1.3 Bounding the operators from above

In order to show the results of existence, we determine that we can set $\{C_k^V\}_{k=1}^K$ and $\{C_k^m\}_{k=1}^K$ recursively as follows:

$$C_1^V = 2K \qquad C_k^V = 2C_{k-1}^V + 2K, \qquad k \in \{2, \dots, K\}$$

$$C_K^m = TK2^{K+5} \qquad C_k^m = TC_{k+1}^m + TK2^{K+5}, \quad k \in \mathcal{K}_{-1}.$$

We observe that for all $k \in \mathcal{K}$, we have $C_k^V \leq C_K^V$ and $C_k^m \leq C_0^m$. From the definition of $\phi^{(0)}$ (C.1), we have

$$\|\phi^{(0)}\|_{\infty} \le \frac{1}{2} \Big(\sum_{k=1}^{K} \|m_k^{(0)}\|_{\infty} \|\Delta V_k^{(0)}\|_{\infty} + \|\eta^{(0)}\|_{\infty} \Big) \le K.$$

In order to bound $\phi^{(1)}$, we take $\epsilon > 0$ small enough such that

$$\begin{split} \|\phi^{(1)}\|_{\infty} &\leq \left\| \frac{1}{2 + \epsilon((\eta^{(0)} + \epsilon \eta^{(1)}))} \right\|_{\infty} \left(\|\eta^{(0)} + \epsilon \eta^{(1)}\|_{\infty} \|\phi^{(0)}\|_{\infty} + \sum_{k=1}^{K} \|m_{k}^{(1)}\|_{\infty} \|\Delta V_{k}^{(0)}\|_{\infty} + \sum_{k=1}^{K} \|m_{k}^{(0)}\|_{\infty} \|\Delta V_{k}^{(1)}\|_{\infty} \\ &+ \epsilon \sum_{k=1}^{K} \|m_{k}^{(1)}\|_{\infty} \|\Delta V_{k}^{(1)}\|_{\infty} + \|\eta^{(1)}\|_{\infty} \right) \\ &\leq 2KC_{0}^{m} + 4KC_{K}^{V}, \end{split}$$

where we used the facts that $0 \leq \eta^{(0)}(t) \leq 1$ for all $t \in [0,T]$ and $0 \leq m_k^{(0)}(t) \leq 1$ for all $k \in \mathcal{K}, t \in [0,T]$ (proved in Appendix B). Similarly, for $\{\lambda_k^{(1)}\}_{k=1}^K$ we get

$$\|\lambda_k^{(1)}\|_{\infty} \le \frac{K}{2} + C_K^V + \frac{\epsilon}{2} \|\phi^{(1)}\|_{\infty}.$$

Consequently,

$$\begin{split} \|S_{k}(V^{(1)}, m^{(1)})\|_{\infty} &\leq T \Big(\|\lambda_{k+1}^{(0)}\|_{\infty} \|m_{k+1}^{(1)}\|_{\infty} + \|m_{k+1}^{(0)}\|_{\infty} \|\lambda_{k+1}^{(1)}\|_{\infty} + \\ \|m_{k}^{(0)}\|_{\infty} \|\lambda_{k}^{(1)}\|_{\infty} + \epsilon \|\Lambda_{k}^{m}\|_{\infty} \Big) \\ &\leq \frac{T}{2} \Big(C_{k+1}^{m} + 2K + 4C_{K}^{V} \Big) + \epsilon T \Big(\|\phi^{(1)}\|_{\infty} + \|\Lambda_{k}^{m}\|_{\infty} \Big), \end{split}$$

where we used the bound $\|\lambda_k^{(0)}\|_{\infty} \leq 1/2$ (Appendix B). As Λ_k^m can be bounded by $\|\Lambda_k^m\|_{\infty} \leq 2C_0^m(C_K^V + \frac{\epsilon}{2} \|\phi^{(1)}\|_{\infty})$, we can take $\epsilon > 0$ small enough such that for all $k \in \mathcal{K}_{-1}$,

$$\epsilon(\|\phi^{(1)}\|_{\infty} + \|\Lambda_k^m\|_{\infty}) \le \frac{1}{2}(C_{k+1}^m + 2K + 4C_K^V).$$

Then we get, for $k \in \mathcal{K}$,

$$||S_k(V^{(1)}, m^{(1)})||_{\infty} \le T\left(C_{k+1}^m + 2K + 4C_K^V\right) \le C_k^m,$$

by observing that, for example, $\|\Lambda_K^m\|_{\infty}$ can be bounded above with the same constant we use for $\|\Lambda_{K-1}^m\|_{\infty}$ to conclude for the case when k = K. This is what we wanted.

Similarly, for integral operators $\{R_k\}_{k=1}^K$ (C.2) related to HJB equations, for sufficiently small $\epsilon > 0$ we have

$$\begin{aligned} \|R_k(V^{(1)}, m^{(1)})\|_{\infty} &\leq \|\phi^{(0)} + V_{k-1}^{(1)}\|_{\infty} + \epsilon \left\| \frac{1}{\lambda_k^{(0)}} \right\|_{\infty} \|\Lambda_k^V\|_{\infty} \\ &\leq \frac{C_k^V}{2} + \epsilon \left\| \frac{1}{\lambda_k^{(0)}} \right\|_{\infty} \|\Lambda_k^V\|_{\infty} \leq C_k^V, \end{aligned}$$

using the fact (proved in Appendix B) that $\max_{k \in \mathcal{K}} \left\| \frac{1}{\lambda_k^{(0)}} \right\|_{\infty} < \infty.$

C.1.4 Applying Schauder Fixed Point Theorem

Now, we have $\Gamma(M) \subset M$. The continuity of $\Gamma : M \to M$ is clear. For compactness, observe that the integrands in R_k (C.2) and S_k (C.3) can be uniformly bounded. Namely, we have a constant C that depends only on the external parameters T, K, r such that for all $(V^{(1)}, m^{(1)}) \in M$, for all $t, s \in [0, 1]$ and for all $k \in \mathcal{K}$, we have

$$|R_k(V^{(1)}, m^{(1)})(t) - R_k(V^{(1)}, m^{(1)})(s)| \le C|t - s|$$

|S_k(V^{(1)}, m^{(1)})(t) - S_k(V^{(1)}, m^{(1)})(s)| \le C|t - s|.

This shows that the family of functions $\Gamma(M)$ is uniformly equicontinuous. Since we are using a uniform norm on our Banach space, the family $\operatorname{clo}(\Gamma(M))$ is also a uniformly equicontinuous family, and since M is a closed set, we have $\operatorname{clo}(\Gamma(M)) \subset M$. Hence $\operatorname{clo}(\Gamma(M))$ is uniformly bounded. Consequently, by the d-dimensional Arzela-Ascoli Theorem, $\operatorname{clo}(\Gamma(M))$ is a compact set. Then by the Schauder Fixed Point Theorem, we conclude that there exists $(V^{(1)}, m^{(1)}) \in M$ such that for all $t \in [0, T]$

$$V_k^{(1)}(t) = \int_t^T e^{-\int_t^s (\lambda_k^{(0)} + r)ds'} \left(\lambda_k^0(\phi^{(0)} + V_{k-1}^{(1)}) + \epsilon \Lambda_k^V(V^{(1)}, m^{(1)})\right) ds, \quad k \in \mathcal{K}$$
(C.4)

and

$$m_{k}^{(1)}(t) = \int_{0}^{t} e^{-\int_{s}^{t} \lambda_{k}^{(0)}(s')ds'} \left(\lambda_{k+1}^{(0)}m_{k+1}^{(1)} - \left(m_{k+1}^{(0)}\lambda_{k+1}^{(1)} - m_{k}^{(0)}\lambda_{k}^{(1)}\right) + \epsilon\Lambda_{k}^{m}(V^{(1)}, m^{(1)})\right)ds, \quad k \in \mathcal{K}_{-1}$$

$$m_{K}^{(1)}(t) = \int_{0}^{t} e^{-\int_{s}^{t} \lambda_{k}^{(0)}(s')ds'} \left(m_{K}^{(0)}\lambda_{K}^{(1)} + \epsilon\Lambda_{K}^{m}(V^{(1)}, m^{(1)})\right)ds.$$
(C.5)

Differentiating both sides of (C.4) and (C.5) with respect to t, we conclude that

$$V_k(t) := V_k^{(0)}(t) + \epsilon V_k^{(1)}(t), \quad k \in \mathcal{K}$$

$$m_k(t) := m_k^{(0)}(t) + \epsilon m_k^{(1)}(t), \quad k \in \mathcal{K}$$

is a solution to our set of coupled ordinary differential equations (2.5).

C.2 **Proof of Uniqueness**

It is important to note that the Schauder Fixed Point Theorem does not guarantee uniqueness. In fact, establishing a priori uniqueness of a solution is not evident from the outset in the context of economic considerations. Nevertheless, when the interactions between the firms are sufficiently small, we can indeed prove uniqueness.

C.2.1 Setup and result for when $\epsilon = 0$

We shall use an energy identity to prove that the solution is unique. To begin, suppose that $(V^{[1]}, m^{[1]})$ and $(V^{[2]}, m^{[2]})$ are two solutions to the equations:

$$\frac{dV_k}{dt} - rV_k + H_k = 0, \quad k \in \mathcal{K}$$

$$\frac{dm_k}{dt} - (G_{k+1}m_{k+1} - G_km_k) = 0, \quad k \in \mathcal{K}$$
(C.6)

where, for $k \in \mathcal{K}$, we defined

$$G_k = \frac{1}{2} \left(1 - \Delta V_k + \epsilon \phi_\epsilon \right) \quad H_k = G_k^2 = \frac{1}{4} \left(1 - \Delta V_k + \epsilon \phi_\epsilon \right)^2,$$

with the understanding that $G_{K+1} = 0$. For the sake of notational brevity, let us denote $\delta L := L^{[2]} - L^{[1]}$ for any generic quantity L. Applying δ to (C.6), multiplying the first equation by δm_k and the second by δV_k , and adding them yields

$$2\frac{d\delta m_k \delta V_k}{dt} - r\delta m_k \delta V_k + \delta m_k \delta H_k - \delta V_k (\delta (G_{k+1}m_{k+1}) - \delta (G_k m_k)) = 0.$$
(C.7)

Since the two solutions agree on the initial and terminal conditions, we have

$$\delta V_k(T) = 0, \quad \delta m_k(0) = 0, \quad k \in \mathcal{K}.$$

We integrate (C.7) and sum over all $k \in \mathcal{K}$. Then, by discrete integration-by-parts, we get

$$\int_0^T e^{-rt} \sum_{k=1}^K \left\{ \delta m_k \delta H_k + \delta (\Delta V_k) \delta (G_k m_k) \right\} dt = 0.$$
 (C.8)

We will now individually target the terms. To simplify this expression, let us introduce another notation: For any generic quantity L, denote $\overline{L} := (L^{[1]} + L^{[2]})/2$. We have the identity $\delta(LF) = \overline{F}\delta L + \overline{L}\delta F$ for another generic quantity F. Using the definitions for H_k and G_k yields $\forall k \in \mathcal{K}$,

$$\delta H_k = 2\overline{G_k}\delta G_k = -\frac{1}{2}\delta(\Delta V_k)\left(1 - \overline{\Delta V_k}\right) + \epsilon Z_k$$
$$\delta(G_k m_k) = \frac{1}{2}\left(\delta m_k(1 - \overline{\Delta V_k}) - \delta(\Delta V_k)\overline{m_k}\right) + \epsilon \zeta_k,$$

where we defined

$$Z_k = -\frac{1}{2} \Big(\overline{\phi_{\epsilon}} \delta(\Delta V_k) - \delta \phi_{\epsilon} (1 - \overline{\Delta V_k}) \Big) + \frac{\epsilon \overline{\phi_{\epsilon}}}{2} \delta \phi_{\epsilon}, \quad \zeta_k = \frac{1}{2} \Big(\overline{\phi_{\epsilon}} \delta m_k + \overline{m_k} \delta \phi_{\epsilon} \Big).$$

Then, (C.8) becomes

$$\frac{1}{2}\int_0^T e^{-rt}\sum_{k=1}^K \overline{m_k}(\delta(\Delta V_k))^2 dt = \epsilon \int_0^T e^{-rt}\sum_{k=1}^K \left\{ Z_k \delta m_k + \zeta_k \delta(\Delta V_k) \right\} dt.$$
(C.9)

Notice that for $\epsilon = 0$, we can conclude that $V^{[1]} = V^{[2]}$ by non-negativity of the left-hand-side integrand. Then, after applying δ to second line in (C.6):

$$\frac{d\delta m_k}{dt} - \left(\delta(G_{k+1}m_{k+1}) - \delta(G_km_k)\right) = 0, \quad k \in \mathcal{K},$$
(C.10)

we can observe that $m^{[1]} = m^{[2]}$ as well, when $\epsilon = 0$.

C.2.2 Uniqueness of V in the general case

For the general case when $\epsilon > 0$, we need to compute the expression on the right-hand side of (C.9) explicitly. To begin, let us observe that

$$Z_k \delta m_k + \zeta_k \delta(\Delta V_k) = \frac{\delta \phi_\epsilon}{2} \Big(\overline{m_k} \delta(\Delta V_k) + \delta m_k (1 - \overline{\Delta V_k}) \Big) + \frac{\epsilon \overline{\phi_\epsilon}}{2} \delta \phi_\epsilon \delta m_k.$$

Therefore,

$$\sum_{k=1}^{K} Z_k \delta m_k + \zeta_k \delta(\Delta V_k) = \frac{1}{2} \delta \phi_\epsilon \sum_{k=1}^{K} \left(\overline{m_k} \delta(\Delta V_k) + \delta m_k (1 - \overline{\Delta V_k}) \right) + \frac{\epsilon \overline{\phi_\epsilon}}{2} \delta \phi_\epsilon \delta \eta.$$
(C.11)

Now let us find $\delta \phi_{\epsilon}$. Recall that we have

$$(2+\epsilon\eta)\phi_{\epsilon} = \sum_{k=1}^{K} m_k (1-\Delta V_k).$$

Applying the difference operator δ on both sides, we have

$$(2+\epsilon\overline{\eta})\delta\phi_{\epsilon}+\epsilon\overline{\phi_{\epsilon}}\delta\eta=\sum_{k=1}^{K}\left(\overline{m_{k}}\delta(\Delta V_{k})-\delta m_{k}(1-\overline{\Delta V_{k}})\right)$$

and thus,

$$\delta\phi_{\epsilon} = \frac{1}{2 + \epsilon\overline{\eta}} \left[\sum_{k=1}^{K} \left(\overline{m_k} \delta(\Delta V_k) - \delta m_k (1 - \overline{\Delta V_k}) \right) - \epsilon \overline{\phi_{\epsilon}} \delta\eta \right].$$
(C.12)

Substituting $\delta \phi_{\epsilon}$ from (C.12) into the right-hand side of (C.11) transforms the equation to

$$\sum_{k=1}^{K} Z_k \delta m_k + \zeta_k \delta(\Delta V_k)$$

$$= \frac{1}{2(2+\epsilon\overline{\eta})} \Big(\sum_{k=1}^{K} \overline{m_k} \delta(\Delta V_k) \Big)^2 - \frac{1}{2(2+\epsilon\overline{\eta})} \Big[\Big(\sum_{k=1}^{K} \delta m_k (1-\overline{\Delta V_k}) \Big) + \epsilon \overline{\phi_\epsilon} \delta \eta \Big]^2 \qquad (C.13)$$

$$\leq \frac{K}{2(2+\epsilon)} \sum_{k=1}^{K} \overline{m_k} (\delta(\Delta V_k))^2,$$

where in the last line we used that $\overline{m_k}(t) \in [0, 1]$ for all $k \in \mathcal{K}$, $t \in [0, T]$ and $\overline{\eta}(t) \in [0, 1]$ for all $t \in [0, T]$. Now combining (C.9) and (C.13), we obtain

$$\frac{1}{2} \int_0^T e^{-rt} \sum_{k=1}^K \overline{m_k} (\delta(\Delta V_k))^2 dt \le \frac{\epsilon K}{2(2+\epsilon)} \int_0^T e^{-rt} \sum_{k=1}^K \overline{m_k} (\delta(\Delta V_k))^2 dt.$$

Consequently, for a sufficiently small $\epsilon > 0$, we get

$$\int_0^T e^{-rt} \sum_{k=1}^K \overline{m_k} (\delta(\Delta V_k))^2 dt = 0.$$

Since $\overline{m_k}(t) > 0$ for $t \in (0,T]$, $k \in \mathcal{K}$, we get $\delta(\Delta V_k(t))^2 = 0$ for all $t \in (0,T]$. From continuity of $V_k(t)$ in t, we have $\delta(\Delta V_k)(0) = 0$ as well. Therefore

$$V^{[1]}(t) = V^{[2]}(t) \quad \forall t \in [0, T].$$

C.2.3 Uniqueness of m in the general case

To see that $m^{[1]} = m^{[2]}$, observe that the second term in (C.13) is identically zero.

$$\frac{1}{2(2+\epsilon\overline{\eta})} \left[\left(\sum_{k=1}^{K} \delta m_k (1-\overline{\Delta V_k}) \right) + \epsilon \overline{\phi_\epsilon} \delta \eta \right]^2 = 0.$$
 (C.14)

With uniqueness of V in mind, combining (C.14) with (C.12) results in $\delta \phi_{\epsilon} = 0$. Therefore, $\phi_{\epsilon}^{[1]}(t) = \phi_{\epsilon}^{[2]}(t)$ for all $t \in [0, T]$. Consequently, the equations in (C.10) become

$$\frac{d\delta m_k}{dt} = -G_k \delta m_k + G_{k+1} \delta m_{k+1}, \quad k \in \mathcal{K}_{-1}$$
$$\frac{d\delta m_K}{dt} = -G_K \delta m_K.$$

Since $\delta m_K(0) = 0$, the solution when k = K is $\delta m_K(t) = 0$ for all $t \in [0, T]$. By backward induction, we find that for any $k \in \mathcal{K}$, we have $\delta m_k(t) = 0$ for all $t \in [0, T]$. Therefore,

$$m^{[1]}(t) = m^{[2]}(t) \quad \forall t \in [0, T].$$

This concludes the proof of uniqueness.

References

- [1] Peter Belobaba. Air Travel Demand and Airline Seat Inventory Management. Ph.D. Dissertation, Massachusetts Institute of Technology, 1987.
- [2] Peter Belobaba. Application of a Probabilistic Decision Model to Airline Seat Inventory Control. Operations Research, 37(2):183–197, 1989.
- [3] Jose M. Betancourt, Ali Hortaçsu, Aniko Oery, and Kevin R. Williams. Dynamic Price Competition: Theory and Evidence from Airline Markets. *National Bureau of Economic Research*, 2022.
- [4] Nicolas Bondoux, Anh Quan Nguyen, Thomas Fiig, and Rodrigo Acuna-Agost. Reinforcement Learning Applied to Airline Revenue Management. Journal of Revenue & Pricing Management, 19:332–348, 2020.
- [5] Pierre Brémaud. Point Processes and Queues: Martingale Dynamics. Springer, 1981.
- [6] René Carmona and François Delarue. Probabilistic Theory of Mean Field Games with Applications I-II. Springer, 2018.
- [7] Patrick Chan and Ronnie Sircar. Bertrand and Cournot Mean Field Games. Applied Mathematics & Optimization, 71:533–569, 2015.
- [8] Patrick Chan and Ronnie Sircar. Fracking, Renewables & Mean Field Games. SIAM Review, 59(3):588–615, 2017.
- [9] Andrew Collins and Lyn Thomas. Comparing Reinforcement Learning Approaches for Solving Game Theoretic Models: A Dynamic Airline Pricing Game Example. *Journal* of Revenue and Pricing Management, 11(1):53–68, 2012.
- [10] Arnoud V. den Boer. Dynamic Pricing and Learning: Historical Origins, Current Research, and New Directions. Surveys in Operations Research and Management Science, 20(1):1–18, 2015.
- [11] Ryan Donnelly and Zi Li. Dynamic Inventory Management with Mean-Field Competition. https://arxiv.org/pdf/2210.17208, 2022. arXiv preprint.
- [12] Wedad Elmaghraby and Pinar Keskinocak. Dynamic Pricing in the Presence of Inventory Considerations: Research Overview, Current Practices, and Future Directions. *Management Science*, 49(10):1287–1309, 2003.
- [13] Guanxing Fu, Paulwin Graewe, Ulrich Horst, and Alexandre Popier. A Mean Field Game of Optimal Portfolio Liquidation. *Mathematics of Operations Research*, 46(4): 1250–1281, 2021. doi: 10.1287/moor.2020.1091.
- [14] Guillermo Gallego and Ming Hu. Dynamic Pricing of Perishable Assets Under Competition. Management Science, 60(5):1241–1259, 2014.

- [15] Guillermo Gallego and Garrett van Ryzin. Optimal Dynamic Pricing of Inventories with Stochastic Demand Over Finite Horizons. *Management Science*, 40(8):999–1020, 1994.
- [16] Nicolás García, Ronnie Sircar, and Mete Soner. Mean Field Games of Control and Cryptocurrency Mining. Submitted, April 2025.
- [17] David Gilbarg and Neil S. Trudinger. Elliptic Partial Differential Equations of Second Order. Springer-Verlag, Berlin, 2nd edition, 2001.
- [18] P. Jameson Graber and Ronnie Sircar. Master equation for Cournot mean field games of control with absorption. *Journal of Differential Equations*, 343:816–909, 2023.
- [19] Paulwin Graewe, Ulrich Horst, and Ronnie Sircar. A Maximum Principle Approach to Deterministic Mean Field Games of Control with Absorption. SIAM Journal on Control and Optimization, 60(5):3173–3190, 2022.
- [20] Fabian Lange, Leonard Dreessen, and Rainer Schlosser. Reinforcement learning versus data-driven dynamic programming: a comparison for finite horizon dynamic pricing markets. *Journal of Revenue and Pricing Management*, 2025.
- [21] Andrew Ledvina. Differential Games in Finance and Economics. Ph.D. Dissertation, Princeton University, Department of Operations Research & Financial Engineering, 2011.
- [22] Zongxi Li, A. Max Reppen, and Ronnie Sircar. A Mean Field Games Model for Cryptocurrency Mining. *Management Science*, 70:2188–2208, 2024.
- [23] Mike Ludkovski and Ronnie Sircar. Exploration and Exhaustibility in Dynamic Cournot Games. *European Journal of Applied Mathematics*, 23(3):343–372, 2012.
- [24] Jeffrey I. McGill and Garrett J. van Ryzin. Revenue Management: Research Overview and Prospects. *Transportation Science*, 33(2):233–256, 1999.
- [25] Kalyan Talluri and Garrett van Ryzin. Revenue Management Under a General Discrete Choice Model of Consumer Behavior. *Management Science*, 50(1):15–33, 2004.
- [26] Xavier Vives. Oligopoly Pricing: Old Ideas and New Tools. MIT Press, 2001.
- [27] Dillon Wong. Flight Prices Dataset. Kaggle Dataset, 2021. URL https://www.kaggle. com/datasets/dilwong/flightprices. Accessed: 2025-04-10.
- [28] Wen Zhao and Yu-Sheng Zheng. Optimal Dynamic Pricing for Perishable Assets with Nonhomogeneous Demand. *Management Science*, 46(3):375–388, 2000.